

With
CITRONELLA

Protects Your Animals

FLYRID[®] PLUS

Description:

Fly-Rid[®] Plus is a multi-species, multi-purpose insect control spray that provides powerful yet safe protection against harmful pests. This ready-to-use permethrin product is for use directly on animals (horses, dogs, goats, sheep, hogs, beef and dairy cattle), their premises (barns, stables, pet beds, porches, poultry premise sites, garages, etc.) as well as other indoor/outdoor applications.

For Use On:

Horses, dogs, goats, sheep, hogs, beef and dairy cattle and their premises, such as barns, stables, pet beds, porches, poultry premise sites and garages. Also approved for other indoor-outdoor applications as indicated by label directions.

Product Advantages:

- Multi-species: For use on Horses, Dogs, Goats, Sheep, Hogs, Beef and Dairy Cattle
- With citronella
- Protects horses and livestock from:
 - Horse flies, horn flies, face flies, house flies, stable flies, deer flies, mosquitoes, and gnats
- Controls fleas and ticks on dogs for 35 days
- Controls fleas (adult and larva) and ticks
- Great for barns, stables and other premises:
 - Controls ticks, which may carry lyme disease (deer tick, lone star tick, brown dog tick, American dog tick and western black legged tick).
 - Provides 60-day control of cockroaches, spiders, ants, crickets, scorpions, millipedes and centipedes.
 - Controls bed bugs and mites in poultry houses.
- Multi-purpose spray
- Permethrin is one of the longer lasting chemicals used in insect repellents
- Ready-to-use
- Convenient, easy-grip bottle
- Powerful protection that's safe, effective and economical



Active Ingredients:

Permethrin.....0.5%

Packaging:

32 oz. spray bottle, 12 bottles per case
UPC# 7-45801-31034-6

Gallon bottle, 4 bottles per case
UPC# 7-45801-31036-0



MADE IN THE USA

040518

FlyRID[®] PLUS

| Tel 800-821-5570 | Fax 816-224-3080 | info@durvet.com |

Protects Your Animals



With
CITRONELLA

Comparison:

COMPARISON BASED ON ACTUAL LABEL CLAIMS	FLY-RID® PLUS Durvet, Inc.	BRONCO® Farnam Companies, Inc.
Active Ingredient	Permethrin.....0.500%	Prallethrin.....0.033% Permethrin.....0.100% Piperonyl Butoxide, Technical.....0.500%
Total Active0.500%0.630%
MUTLI-SPECIES LABEL	YES	NO
PROTECTS HORSES: Controls 6 Fly Species, Gnats, Mosquitoes, & Lice	YES	YES
PROTECTS DOGS 35 DAY CONTROL: Fleas & Ticks on Dogs	YES 35-day control	NO
Fleas (adults and larva) & Ticks	YES	NO
PROTECTS GOATS, SHEEP, CATTLE, & HOGS: Controls 6 Fly Species, Gnats, Mosquitoes, & Lice	YES	NO
PROTECTS POULTRY: Controls Poultry Lice*	YES *Apply only to poultry premise sites	NO
BROAD SPECTRUM TICK CONTROL: Controls 5 types of ticks - including ticks which may carry Lyme Disease. Deer Tick, Lone Star Tick, American Dog Tick, Western Black Legged Tick, Brown Dog Tick.	YES	Only ticks & deer ticks listed
SURFACE SPRAY - 60 DAY CONTROL: Cockroaches, Spiders, Ants, Crickets, & More	YES 60-day control	NO control period listed
CITRONELLA SCENT	YES	YES

All trademarks are property of their respective owners.



040518

FlyRID® PLUS

| Tel 800-821-5570 | Fax 816-224-3080 | info@durvet.com |

Controls 5 species of ticks including those that may carry Lyme Disease.

IF ON SKIN OR ON CLOTHING:	FIRST AID
<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice. 	
Have product container or label with you when calling a poison control center, or doctor, or going in for treatment. For 24-hour emergency information on this product, call 1-800-498-5743 (US & Canada) or 1-857-523-0318 (all other areas).	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

- Long-sleeved shirt and long pants
- Gloves and socks
- Chemical-resistant gloves such as nitrile laminate, nitrile rubber, neoprene rubber or Viton

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been decontaminated or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into streams, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 72 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This pesticide is highly toxic to bees and other pollinating insects exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees or other pollinating insects are foraging the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions, use restrictions and precautions.

SHAKE WELL BEFORE USE

USE RESTRICTIONS

Except when applying to pests or livestock, do not enter or allow others to enter until sprays have dried. Do not use on crops used for food, forage or pasture. Do not apply this product with a sponge. Except when applying to pests and livestock, do not apply this product in a way that will contact workers or other persons, either directly or through drift. Except when applying to pests and livestock, only protective handlers may be in the area during application. When used in Dairy Barns or facilities, Close milk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking. Not for use in outdoor residential misting systems. Do not use in commercial food/feed handling establishments, restaurants or other commercial facilities where food is commercially prepared, processed, served or stored. Do not use in institutions, schools, office buildings or other commercial buildings. Do not breathe vapors or spray mist. Cover or remove all food processing surfaces and utensils during treatment, and thoroughly wash before use. Do not apply when food or drinking water is present. Mix for broadcast use on indoor residential surfaces. Do not use in aircraft cabins. Mix for use in commercial animal premises and tomatic misting systems. Do not contaminate food or feedstuffs.

INDOOR APPLICATION

For spray application, do not use as a space spray indoors. In the home, cover all food processing surfaces and utensils during treatment or thoroughly wash before use. Cover or remove exposed spray nozzles. Remove pets, birds and cover fish aquariums before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry. Do not enter or allow other to enter until sprays have dried. Sulfides staining: Hold sprayer approximately 12 to 15 inches away from surfaces to be treated and spray until surface is damp, but not to the point of run-off; for 60-day control of cockroaches, scorpions, millipedes, centipedes, spiders, ants, crickets. Direct the spray into hiding places, cracks and



FLYRID PLUS



Multi-Purpose Spray

Controls Flies (6 types), Mosquitoes & Gnats on Horses
35-Day Control of Fleas & Ticks on Dogs
 With CITRONELLA

KEEP OUT OF REACH OF CHILDREN	ACTIVE INGREDIENT:	0.5%
CAUTION	PERMETHRIN *	99.5%
See side panels for additional Precautionary Statements	OTHER INGREDIENTS:	100.0%
	TOTAL	100.0%
	*3-(p-henoxphenyl) methyl (1+)-cis-trans-3-(2,4-dichlorophenyl)-2,4-dimethylcyclopropane-carboxylate (Cis/Trans ratio: Min. 55% (1+)-cis and max. 65% (1+)-trans.)	

NET CONTENTS: 1 Qt. (946 mL)

crickets, and behind stallies and drawers. In Control Area: Spray trails, nests and joints of entry.

To Control Fleas Adults and Larvae and Ticks: Including Ticks which may carry Lyme Disease: Adults and Larvae: Spot spray treatment of infested areas, pet beds, resting quarters, kennel crates and crevices, along and behind baseboards, moldings, window and door frames, and for localized treatment of floors and low crevices. Place fresh bedding in animal quarters following treatment. Repeat treatment every 2 weeks for 14 days as needed. Allow dry thoroughly before allowing pets to contact treated areas. To control the source of the infestation, treat your pet with a product registered for animal application. Also, treat your yard with a product registered for yard application.

ON LIVESTOCK: For Use To Protect Cattle, Beef and Dairy Cattle, Goats, Sheep, Hogs, and Horses. In Control of Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head. Do not spray into eyes.

To Control of Stable Flies, Horse Flies, and Deer Flies: Apply at a rate of 7 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at regular intervals to give continued protection.

To Control of Blood Sucking Lice: Apply to the infested areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To Control of Fleas/Lice: Spray nests, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist. To Control of Bed Bugs and Mites: or Fleas/Horses: Spray into crevices of horse racks, cracks in walls, and cracks in nest where the bed bugs and mites hide.

FOR PET DOG USE: Consult a veterinarian before using this product on individualized, medicated, aged, pregnant or nursing animals. Specifications may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

To Control Fleas, Ticks and Lice: Start spraying at the tail/moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and under body. While spraying, lift the hair so that the spray will penetrate into the skin. Make sure spray wets thoroughly, but does not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use protective gloves or mitts when applying this product. Repeat every 15 days as necessary for control.

NOTE: Not for use on puppies less than 12 weeks old. **DO NOT USE ON CATS OR KITTENS.**

OUTDOOR APPLICATION:

Use Restrictions: Do not water the treated area to the point of runoff. Do not make applications during rain. All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: 1) Treatment to soil or vegetation around structures; 2) Applications to lawns, turf, and other vegetation; 3) Applications to building foundations, to a maximum height of 3 feet. Other than applications to building foundations, all outdoor applications to foundations or fences, such as sidewalks, driveways, patios, porches and structural crevices, such as window, doors and eaves) are limited to spot and crack-and-crevice applications only. Application is prohibited directly into sewers or drains, or to any area where a significant discharge to streams, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain or other waterway.

APPLICATION INSTRUCTIONS:

For Use as an External Treatment: In Control: Ticks, including Ticks which may carry Lyme Disease: Use as a residual treatment to control ticks. In Control of Fleas, Lice, Mosquitoes, and Gnats: Use as a residual treatment to control flies, lice, mosquitoes, and gnats. In Control of Fleas, Lice, Mosquitoes, and Gnats: Use as a residual treatment to control flies, lice, mosquitoes, and gnats. In Control of Fleas, Lice, Mosquitoes, and Gnats: Use as a residual treatment to control flies, lice, mosquitoes, and gnats. In Control of Fleas, Lice, Mosquitoes, and Gnats: Use as a residual treatment to control flies, lice, mosquitoes, and gnats.

Do not contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE:** Store in a cool, dry, and secure area. Store in original sealed container away from food or feed. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. If empty, place in trash or other for recycling if available. If partly filled, call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

STORAGE AND DISPOSAL

For questions or comments please call EPA Reg. No. 47000-108-12281
 1-800-821-5570 or email info@duvet.com EPA Est. No. 47002-VA-01
 Manufacturer for Duvet, Inc. EPA Est. No. 61282-VA-01
 100 S. Magellan Drive, Blue Springs, MO 64014 EPA Est. No. 66171-CA-001
 (first letter of batch number indicates establishment) 7 45801 31034 6